

Lie Symmetries and Preliminary Classification of Group-Invariant Solutions of Thomas equation

A. Ouhadan and E. H. El Kinani¹

Université Moulay Ismail, Faculté des Sciences et Techniques, Département de Mathématiques, Laboratoire de Physique Mathématique, Boutalamine B.P.509, Errachidia, Morocco.

UFR de Sciences de l'Ingénieur, Faculté des Sciences et Techniques, Boutalamine B.P.509, Errachidia, Morocco.

Abstract

Using the basic prolongation method and the infinitesimal criterion of invariance, we find the most general Lie point symmetries group of the Thomas equation. Looking the adjoint representation of the obtained symmetry group on its Lie algebra, we will find the preliminary classification of its group-invariant solutions. This latter provides a new exact solutions for the Thomas equation.

Keywords: Lie-point symmetries, Thomas equation, invariant solutions.

2000 Mathematics Subject Classification:70G65, 58K70, 34C14.

¹Junior Associate at The Abdus Salam ICTP, Trieste, Italy. e-mail hkinani@ictp.trieste.it

1 Introduction

The Lie group method is one of the most powerful method available to analyse nonlinear evolution systems and their methods of solution [1, 2, 3, 4, 5]. Introduced firstly by Sophus Lie [6] and consists of a systematic procedure for the determination of continuous symmetry transformations of a system of nonlinear partial differential equations (NLPDE). An important feature of this method is that one can derive special solutions associated with nonlinear PDEs straightforwardly which are otherwise inaccessible through other methods. One of the obvious advantages of knowing a symmetry group of a system of differential equations is that one can use the defining propriety of such a group and construct new solutions to the system from known ones. One can also use symmetry group to explicitly determine types of solution which are themselves invariant under some subgroups of the full symmetry group of the system and provides a new classification of different classes of solutions. The basic idea of the Lie symmetry method is to find the transformations groups associated with a given systems under a continuous group of transformations and to find a reduction transformation from the symmetries. In particular for nonlinear partial differential equations with two independent variables the reduction transformation can be used to reduce the number of independent variables by one. In the case of the ordinary differential equations the determination of a one-parameter symmetry group allows the reduction of the order of the equation by one. Here using the criterion of invariance of the equation under the infinitesimal prolonged generators, we find the most general Lie point symmetries group of the Thomas equation. Looking the adjoint representation of the obtained symmetry group on its Lie algebra, we will find the preliminary classification of group-invariant solutions. This provides a new exact solutions of the Thomas equation. The paper is arranged as follows. In section 2, we introduce the Thomas equation and by using the prolongation formulae and the infinitesimal criterion of invariance we determine the most general symmetry group and the corresponding Lie algebra for Thomas equation. Section 3, is devoted to the construction of the of group-invariant solutions and its classification which provides in each case new exact solutions for the Thomas equation. In section 4, we present conclusions and finally in the Appendix, we briefly give the methods for constructing the invariants of the differential equation.

2 Determination of Symmetry group for Thomas equation

2.1 Thomas equation

The Thomas equation is an interesting subject in physical sciences [7, 8, 9]it arises in the study of chemical exchange processus and sometime called Thompson equation. Proposed by Thomas [9] and has the form:

$$u_{xy} + \alpha u_x + \beta u_y + \gamma u_x u_y = 0, \quad (1)$$

where : $u_x = \frac{\partial u}{\partial x}$, $u_y = \frac{\partial u}{\partial y}$, $u_{xy} = \frac{\partial^2 u}{\partial x \partial y}$ and α, β and γ are constants. For exchange processus $\alpha > 0, \beta > 0$ and $\gamma \neq 0$.

2.2 Lie algebra of Thomas equation

In order to calculate the Lie point symmetries of the system(1), we first consider the generator of the group of point transformations. Here there are two independent variables x and y , and one dependent variable u . The Thomas equation is of second order ($n = 2$). Then a general vector field on the space of independent and dependent variables ($X \times R$ where $X = \langle x, y \rangle$) takes the form :

$$v = \xi(x, y, u) \frac{\partial}{\partial x} + \eta(x, y, u) \frac{\partial}{\partial y} + \varphi(x, y, u) \frac{\partial}{\partial u}. \quad (2)$$

Where ξ, η and φ depend on x, y and u . We wish to determine all possible coefficient functions ξ, η and φ , so that the corresponding one-parameter group $\exp \epsilon v$ is a symmetry group of Thomas equation. According the technics using in [1], since the Thomas equation is of second order, we need to know the second prolongation $Pr^2 v$ of the infinitesimal generators which is given by the following formulae :

$$Pr^2 v = v + \varphi^x \frac{\partial}{\partial u_x} + \varphi^y \frac{\partial}{\partial u_y} + \varphi^{xx} \frac{\partial}{\partial u_{xx}} + \varphi^{xy} \frac{\partial}{\partial u_{xy}} + \varphi^{yy} \frac{\partial}{\partial u_{yy}}, \quad (3)$$

where the coefficients $\varphi^x, \varphi^y, \varphi^{xx}, \varphi^{xy}$ and φ^{yy} are given by :

$$\varphi^x = D_x \varphi - u_x D_x \xi - u_y D_x \eta, \quad (4)$$

$$\varphi^y = D_y \varphi - u_x D_y \xi - u_y D_y \eta, \quad (5)$$

$$\varphi^{xy} = D_{xy}(\varphi - \xi u_x - \eta u_y) + \xi u_{xxy} + \eta u_{xyy}, \quad (6)$$

$$\varphi^{yy} = D_{yy} \varphi - 2u_{xy} D_y \xi - 2u_{yy} D_y \eta - u_x D_{yy} \xi - u_y D_{yy} \eta, \quad (7)$$

$$\varphi^{xx} = D_{xx} \varphi - 2u_{xx} D_x \xi - 2u_{xy} D_x \eta - u_x D_{xx} \xi - u_y D_{xx} \eta, \quad (8)$$

where D_x and D_y are total derivatives on x and y respectively and $D_{xy} = D_x(D_y)$. After a long calculation, we obtain :

$$\varphi^x = \varphi_x + (\varphi_u - \xi_x)u_x - \eta_x u_y - \xi_u u_x^2 - \eta_u u_x u_y \quad (9)$$

$$\varphi^y = \varphi_y + (\varphi_u - \eta_y)u_y - \xi_y u_x - \eta_u u_y^2 - \xi_u u_x u_y \quad (10)$$

$$\varphi^{xy} = \varphi_{xy} + (\varphi_{yu} - \xi_{xy})u_x + (\varphi_{xu} - \eta_{xy})u_y - \xi_{yu} u_x^2 \quad (11)$$

$$+ (\varphi_{uu} - \xi_{ux} - \eta_{yu})u_x u_y - \eta_{ux} u_y^2 - \xi_{uu} u_x^2 u_y - \eta_{uu} u_x u_y^2 \\ - 2\xi_u u_x u_{xy} - 2\eta_u u_y u_{xy} - \xi_y u_{xx} - \eta_x u_{yy} - \xi_u u_y u_{xx} - \eta_u u_x u_{yy} \\ + (\varphi_u - \xi_x - \eta_y)u_{xy}$$

$$\varphi^{xx} = \varphi_{xx} + (2\varphi_{xu} - \xi_{xx})u_x - \eta_{xx} u_y - 2\eta_{xu} u_x u_y \quad (12)$$

$$\begin{aligned}
& + (\varphi_{uu} - 2\xi_{xu})u_x^2 - \eta_{uu}u_x^2u_y - 2\eta_xu_{xy} \\
& - 2\eta_uu_xu_{xy} + (\varphi_u - 2\xi_x)u_{xx} - 3\xi_uu_xu_{xx} - \xi_{uu}u_x^3 \\
& - \eta_uu_yu_{xx} \\
\varphi^{yy} = & \varphi_{yy} + (2\varphi_{yu} - \eta_{yy})u_y - \xi_{yy}u_x - 2\xi_{yu}u_xu_y \\
& + (\varphi_{uu} - 2\eta_{yu})u_y^2 - \xi_{uu}u_y^2u_x - 2\xi_yu_{xy} \\
& - 2\xi_uu_yu_{xy} + (\varphi_u - 2\eta_y)u_{yy} - 3\eta_uu_yu_{yy} - \eta_{uu}u_y^3 \\
& - \xi_uu_xu_{yy}
\end{aligned} \tag{13}$$

To use the infinitesimal criterion of invariance, let us introduce $\Delta = u_{xy} + \alpha u_x + \beta u_y + \gamma u_x u_y$.

Suppose that :

$$\Delta = 0, \quad \text{if } Pr^2v(\Delta) = 0, \tag{14}$$

for every infinitesimal generators v of G , then G is a symmetry group of the Thomas equation (1). Substituting the general formulae (9, 10) and (11) into (3) and (14), after eliminating the dependence between the derivatives coming from the equation itself and equating the coefficients of the various monomials in the first and second order, we find the defining equations for the symmetry group of the Thomas equation to be the following :

Monomial	coefficient
1	$\alpha\varphi_x + \beta\varphi_y + \varphi_{xy}$
u_x	$\varphi_{yu} - \xi_{xy} + \gamma\varphi_y - \beta\xi_y + \alpha\eta_y$
u_y	$\varphi_{xu} - \eta_{xy} + \gamma\varphi_x - \alpha\eta_x + \beta\xi_x$
$u_x u_y$	$\varphi_{uu} - \xi_{ux} - \eta_{yu} + \beta\xi_u + \alpha\eta_u + \gamma\varphi_u$
u_x^2	$-\xi_{yu} - \gamma\xi_y + \alpha\xi_u$
u_y^2	$-\eta_{xu} - \gamma\eta_x + \beta\eta_u$
$u_x^2 u_y$	$-\xi_{uu}$
$u_x u_y^2$	$-\eta_{uu}$
u_{xx}	$-\xi_y$
u_{yy}	$-\eta_x$
$u_y u_{xx}$	$-\xi_u$
$u_x u_{yy}$	$-\eta_u$

Table 1: Coefficients of various monomials

The system determining the symmetry group of our equation is :

$$\alpha\varphi_x + \beta\varphi_y + \varphi_{xy} = 0; \tag{15-1}$$

$$\varphi_{yu} - \xi_{xy} + \gamma\varphi_y - \beta\xi_y + \alpha\eta_y = 0; \tag{15-2}$$

$$\varphi_{xu} - \eta_{xy} + \gamma\varphi_x - \alpha\eta_x + \beta\xi_x = 0; \tag{15-3}$$

$$\varphi_{uu} - \xi_{ux} - \eta_{yu} + \beta\xi_u + \alpha\eta_u + \gamma\varphi_u = 0; \tag{15-4}$$

$$-\xi_{yu} - \gamma\xi_y + \alpha\xi_u = 0; \tag{15-5}$$

$$-\eta_{xu} - \gamma\eta_x + \beta\eta_u = 0; \quad (15-6)$$

$$-\xi_{uu} = 0; \quad (15-7)$$

$$-\eta_{uu} = 0; \quad (15-8)$$

$$-\xi_y = 0; \quad (15-9)$$

$$-\eta_x = 0; \quad (15-10)$$

$$-\xi_u = 0; \quad (15-11)$$

$$-\eta_u = 0; \quad (15-12)$$

From (15-9) to (15-12) we obtain :

$$\xi = \xi(x); \quad \text{and} \quad \eta = \eta(y)$$

From (15-4) we deduce that :

$$\varphi(x, y, u) = g(x, y) \exp^{-\gamma u} + N(x, y)$$

For convenient we can chose $\frac{-g(x,y)}{\gamma}$ for $g(x, y)$.

Using the above results and (15-2) with (15-3) one finds :

$$\gamma N_x + \beta \xi_x = 0; \quad \gamma N_y + \alpha \eta_y = 0,$$

which yields to :

$$N_{xy} = 0; \quad \xi_x = -\frac{\gamma N_x}{\beta}, \quad \text{and} \quad \eta_y = -\frac{\gamma N_y}{\alpha},$$

so we have $N(x, y) = mx + ny + a$ where m, n and a are arbitrary constants, the linear first partial differential equation (15-1) leads to $\alpha m + \beta n = 0$.

Then, for $k = \frac{m}{\beta}$, one obtains the most general form of the coefficient functions ξ, η and φ which are the form :

$$\xi(x, y, u) = -k\gamma x + c, \quad (15)$$

$$\eta(x, y, u) = k\gamma y + b, \quad (16)$$

$$\varphi(x, y, u) = -\frac{g(x, y)}{\gamma} e^{-\gamma u} + k(\beta x - \alpha y) + a, \quad (17)$$

where a, b, c and k are arbitrary constants and $g(x, y)$ an arbitrary solution of the equation (15-1). Then, we obtain the most general form for vector field v which is :

$$v = (-k\gamma x + c) \frac{\partial}{\partial x} + (k\gamma y + b) \frac{\partial}{\partial y} + \left(-\frac{g(x, y)}{\gamma} e^{-\gamma u} + k(\beta x - \alpha y) + a \right) \frac{\partial}{\partial u}. \quad (18)$$

Hence, the Lie algebra of infinitesimal symmetry of the Thomas equation is spanned by the four vectors fields :

$$v_1 = \frac{\partial}{\partial x}; \quad (19)$$

$$v_2 = \frac{\partial}{\partial y}; \quad (20)$$

$$v_3 = \frac{\partial}{\partial u}; \quad (21)$$

$$v_4 = -\gamma x \frac{\partial}{\partial x} + \gamma y \frac{\partial}{\partial y} + (\beta x - \alpha y) \frac{\partial}{\partial u}; \quad (22)$$

and the infinite dimensional subalgebra :

$$v_g = -\frac{g(x, y)}{\gamma} e^{-\gamma u} \frac{\partial}{\partial u}. \quad (23)$$

The one-parameter groups G_i generated by the v_i are given in the following table. The entries gives the transformed point $\exp(\epsilon v_i)(x, y, u) = (\tilde{x}, \tilde{y}, \tilde{u})$:

$$G_1 : (x + \epsilon, y, u) \quad (24)$$

$$G_2 : (x, y + \epsilon, u) \quad (25)$$

$$G_3 : (x, y, u + \epsilon) \quad (26)$$

$$G_4 : (xe^{-\gamma\epsilon}, ye^{\gamma\epsilon}, \frac{\beta}{\gamma}x(1 - e^{-\gamma\epsilon}) + \frac{\alpha}{\gamma}y(1 - e^{\gamma\epsilon}) + u) \quad (27)$$

$$G_g : (x, y, \frac{1}{\gamma} \log[\gamma g(x, y)\epsilon + e^{\gamma u}]) \quad (28)$$

Since each group G_i is a symmetry group, then if $u = f(x, y)$ is a solution of the Thomas equation, so are the functions :

$$u_1 = f(x - \epsilon, y) \quad (29)$$

$$u_2 = f(x, y - \epsilon,) \quad (30)$$

$$u_3 = f(x, y) + \epsilon \quad (31)$$

$$u_4 = \frac{\beta}{\gamma}x(e^{\gamma\epsilon} - 1) + \frac{\alpha}{\gamma}y(e^{-\gamma\epsilon} - 1) + f(xe^{\gamma\epsilon}, ye^{-\gamma\epsilon}) \quad (32)$$

$$u_g = \frac{1}{\gamma} \log[\gamma g(x, y)\epsilon + e^{\gamma f(x, y)}]. \quad (33)$$

where ϵ is any number real.

The commutation relations between these vector fields are given in the following table :

$[v_i, v_j]$	v_1	v_2	v_3	v_4	v_f
v_1	0	0	0	$-\gamma v_1 + \beta v_3$	v_{f_x}
v_2	0	0	0	$\gamma v_2 - \alpha v_3$	v_{f_y}
v_3	0	0	0	0	$v_{-\gamma f}$
v_4	$\gamma v_1 - \beta v_3$	$-\gamma v_2 + \alpha v_3$	0	0	v_ψ
v_g	$-v_{f_x}$	$-v_{f_y}$	$v_{\gamma f}$	v_ψ	0

Table 2: Commutator table for the Lie algebra v_i and v_g

where

$$\psi = -\gamma x f_x + \gamma y f_y - \gamma(\beta x - \alpha y)f.$$

Since the totality of infinitesimal symmetries must be a Lie algebra, we can conclude that if $f(x, y)$ is any solution of the equation in (15-1), so are f_x , f_y and $-\gamma x f_x + \gamma y f_y - \gamma(\beta x - \alpha y)f$.

3 Classification of Group-Invariant Solutions

Recall first that in general to each one parameter subgroups of the full symmetry group of a system there will corresponding a family of solutions, such solutions are called invariant solutions [1]. In this paper we are interesting only on the symmetry algebra g of the Thomas equation which is spanned by the vector fields v_1, v_2, v_3 and v_4 . Every one-dimensional subalgebra of g is determined by a nonzero vector v of the form :

$$v = a_1 v_1 + a_2 v_2 + a_3 v_3 + a_4 v_4, \quad (34)$$

where a_i are arbitrary constants. Our task is to simplify as many of the coefficients a_i as possible through application of adjoint to v . To compute the adjoint representation, we use the Lie series :

$$Ad(\exp(\epsilon v_i))v_j = v_j - \epsilon[v_i, v_j] + \frac{\epsilon^2}{2}[v_i, [v_i, v_j]] - \dots \quad (35)$$

Then from the commutation table 2, we obtain the following table :

Ad	v_1	v_2	v_3	v_4
v_1	v_1	v_2	v_3	$v_4 + \epsilon \gamma v_1 - \epsilon \beta v_3$
v_2	v_1	v_2	v_3	$v_4 - \epsilon \gamma v_2 + \epsilon \alpha v_3$
v_3	v_1	v_2	v_3	v_4
v_4	$e^{-\gamma \epsilon} v_1 + \frac{\beta}{\gamma}(1 - e^{-\gamma \epsilon})v_3$	$e^{\gamma \epsilon} v_2 - \frac{\alpha}{\gamma}(e^{-\gamma \epsilon} - 1)v_3$	v_3	v_4

Table 3: Adjoint table for the Lie algebra v_i

In what follows, we begin the classification process :

Case.1

Suppose first that $a_4 \neq 0$, we can assume that $a_4 = 1$, and the vector field v takes the form :

$$v = a_1 v_1 + a_2 v_2 + a_3 v_3 + v_4 \quad (36)$$

Refereing to the table.3, if we act on such a v by $Ad(\exp(\frac{a_3}{\beta})v)$, then we can make the coefficient v_3 vanish . Then the reduced vector field takes the form :

$$v = (a_1 - \gamma x) \frac{\partial}{\partial x} + (a_2 + \gamma y) \frac{\partial}{\partial y} + (\beta x - \alpha y) \frac{\partial}{\partial u}. \quad (37)$$

The invariants ς and χ can be found by integrating the corresponding characteristic system (see *Appendix for construction of the invariants*), which is :

$$\frac{dx}{a_1 - \gamma x} = \frac{dy}{a_2 + \gamma y} = \frac{du}{\beta x - \alpha y} \quad (38)$$

The obtained solution are given by

$$\chi = (a_1 - \gamma x)(a_2 + \gamma y), \quad (39)$$

and

$$\varsigma = u + \frac{\beta}{\gamma} x + \frac{(\beta a_1 + \alpha a_2)}{\gamma^2} \log(\gamma x - a_1) - \frac{\alpha \chi}{\gamma^2(\gamma x - a_1)}. \quad (40)$$

Therefore, a solution of our equation in this case is :

$$u = f(x, \chi, \varsigma) = \varsigma - \left(\frac{\beta a_1 + \alpha a_2}{\gamma^2} \right) \log(\gamma x - a_1) + \frac{\alpha \chi}{\gamma^2(\gamma x - a_1)^2} - \frac{\beta}{\gamma} x \quad (41)$$

The derivatives of u are given in terms of ς and χ as :

$$u_x = -\gamma(a_2 + \gamma y)\varsigma_\chi - \frac{\beta a_1 + \alpha a_2}{\gamma(\gamma x - a_1)} - \frac{\beta}{\gamma}; \quad (42)$$

$$u_y = -\frac{\alpha}{\gamma} + \gamma(a_1 - \gamma x)\varsigma_\chi; \quad (43)$$

$$u_{xy} = -\gamma^2 \varsigma_\chi - \gamma^2 \varsigma_{\chi\chi}; \quad (44)$$

Substituting (42), (43) and (44) into the Thomas equation we obtain reduced equation of the form :

$$\gamma^2 \chi \varsigma_{\chi\chi} + \gamma^3 \chi \varsigma_\chi^2 + \gamma^2 (\gamma - \beta a_1 - \alpha a_2) \varsigma_\chi + \frac{\alpha \beta}{\gamma} = 0. \quad (45)$$

For $\varsigma_\chi = \theta$, one obtains the following equation :

$$\theta' = -\gamma \theta^2 - \frac{(\gamma - \beta a_1 - \alpha a_2)}{\gamma} \frac{\theta}{\chi} + \frac{\alpha \beta}{\gamma^2 \chi} = 0; \quad (46)$$

this Riccati type equation is transformed by using the change :

$$y(\chi) = \exp\left(\int \gamma \theta(\chi)\right), \quad (47)$$

multiplying by χ one obtain the Fuchs type equation :

$$\chi^2 y'' + e \chi y' + m \chi y = 0, \quad (48)$$

where $e = \frac{(\gamma - \beta a_1 - \alpha a_2)}{\gamma}$ and $m = \frac{\alpha \beta}{\gamma^2}$.

This equation admits a solution developable in series in the neighboring of zero, for $\chi > 0$, we have :

$$y_p(\chi) = \sum_{n=0}^{\infty} a_n \chi^n, \quad (49)$$

where $a_0 = 1$ and $a_n = \frac{(-m)^n}{n!e(e+1)\dots(e+n-1)}$, $n \geq 1$

In the case $\chi < 0$ by introducing the transformation $\chi \mapsto -\chi$, This series have an infinite radius of convergence for all $\chi > 0$, then if we introduce the change $z_p(\chi) = \frac{y_p'}{\gamma y_p}$, z_p is a particular solution of the equation (46). Putting $\theta = z + z_p$, (46) becomes a Bernoulli type equation :

$$z' = -\gamma z^2 - 2\gamma z_p z, \quad (50)$$

which is transformed in the linear equation (with $z = \frac{1}{f}$)

$$f' = 2\gamma z_p f + \gamma. \quad (51)$$

The solution of (51) is :

$$f(\chi) = g_p(\chi) y_p^2, \quad (52)$$

where g_p is a primitive of $\frac{1}{\gamma y_p^2}$

Therefore,

$$\theta(\chi) = \frac{1}{f(\chi)} + z_p(\chi). \quad (53)$$

Thus, we have $\varsigma = \varsigma(\chi) = \int \theta(\chi) d\chi$,

and then, we obtain the solution in this case which is :

$$u = \varsigma(\chi) - \left(\frac{\beta a_1 + \alpha a_2}{\gamma^2}\right) \log(\gamma x - a_1) + \frac{\alpha \chi}{\gamma^2(\gamma x - a_1)} - \frac{\beta}{\gamma} x, \quad (54)$$

where $\chi = (a_1 - \gamma x)(a_2 + \gamma y)$.

Case.2

Suppose now that $a_4 = 0$, and $a_3 \neq 0$. In this situation the invariants associated to the field vector v are of the form :

Case 2.1 : If $a_1 a_2 \neq 0$,

$$\begin{aligned} v &= a_1 v_1 + a_2 v_2 + v_3; \\ &= a_1 \frac{\partial}{\partial x} + a_2 \frac{\partial}{\partial y} + \frac{\partial}{\partial u}; \end{aligned} \quad (55)$$

The invariants are :

$$\chi = a_2 x - a_1 y; \quad (56)$$

$$\varsigma = u - \frac{y}{a_2}; \quad (57)$$

Next, we have :

$$u_x = a_2 \varsigma_\chi; \quad (58)$$

$$u_y = \frac{1}{a_2} - a_1 \varsigma_\chi; \quad (59)$$

$$u_{xy} = -a_1 a_2 \varsigma_{\chi\chi}. \quad (60)$$

Substituting these quantities in the Thomas equation, we find the reduced equation :

$$-a_1 a_2 \varsigma_{\chi\chi} + (\alpha a_2 - \beta a_1 + \gamma) \varsigma_\chi - a_1 a_2 \gamma \varsigma_\chi^2 + \frac{\beta}{a_2} = 0; \quad (61)$$

Putting $\varsigma_\chi = \theta$, then (61) takes the form :

$$-a_1 a_2 \theta' + (\alpha a_2 - \beta a_1 + \gamma) \theta - a_1 a_2 \gamma \theta^2 + \frac{\beta}{a_2} = 0; \quad (62)$$

which is the Riccati type equation.

Case 2.1.a: If $(\alpha a_1 - \beta a_1 + \gamma)^2 + 4a_1 \gamma \beta \geq 0$.

Then, our equation admit one constant solution θ_0 which satisfying the second order equation:

$$(\alpha a_1 - \beta a_1 + \gamma)z - a_1 a_2 \gamma z^2 + \frac{\beta}{a_2} = 0. \quad (63)$$

Then by redefining the variable $\theta = f + \theta_0$, one obtain the following Bernoulli type equation :

$$-a_1 a_2 f' + (\alpha a_2 - \beta a_1 + \gamma - 2a_1 a_2 \gamma \theta_0) f - a_1 a_2 \gamma f^2 = 0; \quad (64)$$

Put now $f = \frac{1}{h}$ one obtain a linear equation:

$$h' = \frac{2a_1a_2\gamma\theta_0 - \alpha a_2 + \beta a_1 + \gamma}{a_1a_2}h + \gamma \quad (65)$$

This equation admit a solution, since $\frac{2a_1a_2\gamma\theta_0 - \alpha a_2 + \beta a_1 + \gamma}{a_1a_2} \neq 0$, which is :

$$h = A \exp\left(\frac{2a_1a_2\gamma\theta_0 - \alpha a_2 - \beta a_1 + \gamma}{a_1a_2}\chi\right) - \frac{\gamma a_1a_2}{2a_1a_2\gamma\theta_0 - \alpha a_1 - \beta a_1 + \gamma} \quad (66)$$

where A is an arbitrary constant. Hence,

$$\varsigma = \int \frac{1}{A \exp\left(\frac{2a_1a_2\gamma\theta_0 - \alpha a_2 + \beta a_1 + \gamma}{a_1a_2}\chi\right) - \frac{\gamma a_1a_2}{2a_1a_2\gamma\theta_0 - \alpha a_1 - \beta a_1 + \gamma}} d\chi + \theta_0\chi + cte \quad (67)$$

After integration we obtain :

$$\varsigma = \frac{1}{\gamma} \log\left(A - \frac{\gamma}{C} e^{-C\chi}\right) + \theta_0\chi + cte \quad (68)$$

where $C = \frac{2a_1a_2\gamma\theta_0 - \alpha a_2 + \beta a_1 - \gamma}{a_1a_2}$. Finally, the solution of the Thomas equation is:

$$u = \frac{1}{\gamma} \log\left(A - \frac{\gamma}{C} e^{-C(a_2x - a_1y)}\right) + \theta_0(a_2x - a_1y) + \frac{y}{a_2} + cte; \quad (69)$$

Case 2.1.b: If $(\alpha a_2 - \beta a_1 + \gamma)^2 + 4a_1\gamma\beta < 0$.

In this situation we have :

$$\frac{(\alpha a_2 - \beta a_1 + \gamma)}{a_1a_2}\theta - \gamma\theta^2 + \frac{\beta}{a_1a_2^2} \neq 0. \quad (70)$$

The reduced equation obtained can be written in the separation form as :

$$\frac{d\theta}{\frac{(\alpha a_2 - \beta a_1 + \gamma)}{a_1a_2}\theta - \gamma\theta^2 + \frac{\beta}{a_1a_2^2}} = dx. \quad (71)$$

If we put $A_1 = \frac{(\alpha a_2 - \beta a_1 + \gamma)}{a_1a_2}$, $A_2 = -\gamma$, and $A_3 = \frac{\beta}{a_1a_2^2}$, thus the problem becomes to the integration of the following equation :

$$\frac{d\theta}{A_1\theta + A_2\theta^2 + A_3} = dx. \quad (72)$$

Let $\Xi = \frac{4A_2A_3 - A_1^2}{4A_2^2}$, $\Xi > 0$. Then, we look to the integration of

$$\frac{1}{(\theta + \frac{A_1}{2A_2})^2 + \Xi}. \quad (73)$$

By using the change of the variable $t = \Xi + \frac{A_1}{2A_2}$, we have :

$$\int \frac{d\theta}{(\theta + \frac{A_1}{2A_2})^2 + \Xi} = \frac{1}{\sqrt{\Xi}} \arctan\left(\frac{\theta + \frac{A_1}{2A_2}}{\sqrt{\Xi}}\right) + cte; \quad (74)$$

and therefore,

$$\frac{1}{\sqrt{\Xi}} \arctan\left(\frac{\theta + \frac{A_1}{2A_2}}{\sqrt{\Xi}}\right) = A_2 x + cte, \quad (75)$$

then,

$$\theta = \sqrt{\Xi} tg(A_2 \sqrt{\Xi} x + cte) - \frac{A_1}{2A_2}, \quad (76)$$

and we obtain :

$$\varsigma_\chi = \sqrt{\Xi} tg(A_2 \sqrt{\Xi} \chi + cte) - \frac{A_1}{2A_2}, \quad (77)$$

$$\begin{aligned} \varsigma &= \sqrt{\Xi} \int tg(A_2 \sqrt{\Xi} \chi + cte) d\chi - \frac{A_1}{2A_2} \chi + cte, \\ &= \frac{1}{2A_2} \log(1 + tg(A_2 \sqrt{\Xi} \chi + cte)) - \frac{A_1}{2A_2} \chi + cte. \end{aligned} \quad (78)$$

Next, we find the solution of the Thomas equation which is :

$$u = \frac{1}{2A_1} \log(1 + tg(A_2 \sqrt{\Xi}(a_2 x - a_1 y) + A_0)) - \frac{A_1}{2A_2} (a_2 x - a_1 y) + \frac{y}{a_2} + cte; \quad (79)$$

where $A_0 = \text{constant}$.

Case 2.2 : If $a_2 = 0, a_1 \neq 0$ and $a_1 \neq \frac{-\gamma}{\beta}$

In this case the vectors field is:

$$v = a_1 \frac{\partial}{\partial x} + \frac{\partial}{\partial u}. \quad (80)$$

The invariants are :

$$\chi = y, \quad (81)$$

$$\varsigma = -a_1 u + x, \quad (82)$$

then, $u = \frac{-\chi}{a_1} + \frac{x}{a_1}$, and :

$$u_x = \frac{1}{a_1}, \quad (83)$$

$$u_y = \frac{-\varsigma_\chi}{a_1}, \quad (84)$$

$$u_{xy} = 0. \quad (85)$$

After substituting these expressions into the Thomas equation, we obtain the following reduced equation :

$$(\beta a_1 + \gamma)\varsigma_\chi = a_1\alpha; \quad (86)$$

and then,

$$\varsigma = \frac{a_1\alpha}{\beta a_1 + \gamma}\chi + cte. \quad (87)$$

Next we have,

$$u = \frac{x}{a_1} - \frac{\alpha}{\beta a_1 + \gamma}\varsigma + cte; \quad (88)$$

and therefore,

$$u = \frac{x}{a_1} - \frac{\alpha}{\beta a_1 + \gamma}y + cte. \quad (89)$$

Case 2.3 : If $a_1 = \frac{-\gamma}{\beta}$.

The vector field v take the form :

$$v = \frac{-\gamma}{\beta} \frac{\partial}{\partial x} + \frac{\partial}{\partial u}. \quad (90)$$

and the invariants are :

$$\chi = y; \quad (91)$$

$$\varsigma = \beta x + \gamma u. \quad (92)$$

Therefore

$$u = \frac{\chi}{\gamma} - \frac{\beta}{\gamma}x, \quad (93)$$

and then,

$$u_x = \frac{-\beta}{\gamma}; \quad (94)$$

$$u_y = \frac{-\varsigma_\chi}{\gamma}; \quad (95)$$

$$u_{xy} = 0. \quad (96)$$

Substituting the above expressions into Thomas equation, one find that $\alpha\beta = 0$. Then we conclude that there is not the solution in this case.

Case 2.4 : If $a_1 = a_2 = 0$.

we have :

$$v = \frac{\partial}{\partial u}; \quad (97)$$

the invariants are $\varsigma = x$, and $\chi = y$, then like in the last case there is not the solution of the equation.

Case.3

In this case we choose $a_4 = a_3 = 0$, then the vector field is the form :

$$v = a_1 \frac{\partial}{\partial x} + a_2 \frac{\partial}{\partial y}; \quad (98)$$

Case 3.1 : $a_1 a_2 \neq 0$.

The vector field is of the form :

$$v = \frac{\partial}{\partial x} + a_2 \frac{\partial}{\partial y}; \quad (99)$$

and the invariants ς et χ are solutions of the following characteristics system :

$$\frac{dx}{1} = \frac{dy}{a_2} = \frac{du}{0}. \quad (100)$$

The solutions are :

$$\chi = x - \frac{y}{a_2}; \quad (101)$$

$$\varsigma = u. \quad (102)$$

next we have : $u = \varsigma$, therefore :

$$u_x = \varsigma_\chi; \quad (103)$$

$$u_y = \frac{-\varsigma_\chi}{a_2}; \quad (104)$$

$$u_{xy} = \frac{-\varsigma_{\chi\chi}}{a_2}; \quad (105)$$

By substituting these expressions into the Thomas equation, we obtain the following equation :

$$\varsigma_{\chi\chi} + (\beta - a_2\alpha)\varsigma_\chi + \gamma\varsigma_\chi^2 = 0; \quad (106)$$

putting $\theta = \varsigma_\chi$, we obtain the Bernoulli equation :

$$\theta' + (\beta - a_2\alpha)\theta + \gamma\theta^2 = 0 \quad (107)$$

Case 3.1.a : If $a_2 = \frac{\beta}{\alpha}$.

The reduced equation takes the form :

$$\theta' + \gamma\theta^2 = 0; \quad (108)$$

which have the solution :

$$\theta = \frac{1}{\gamma\chi + cte}, \quad (109)$$

and therefore,

$$u = \frac{1}{\gamma} \log(\gamma(x - \frac{y}{a_2}) + k_0) + cte; \quad (110)$$

where k_0 , is arbitrary constant.

Case 3.1.b : If $a_2 \neq \frac{\beta}{\alpha}$.

We adopt the change $\theta = \frac{1}{z}$, we obtain the following linear equation :

$$z' = (\beta - a_2\alpha)z + \gamma, \quad (111)$$

which admit the general solution :

$$z = ke^{(\beta - a_2\alpha)\chi} - \frac{\gamma}{\beta - a_2\alpha}. \quad (112)$$

Therefore,

$$v = \int \frac{1}{ke^{(\beta - a_2\alpha)\chi} - \frac{\gamma}{\beta - a_2\alpha}} d\chi, \quad (113)$$

and then,

$$u = \frac{-1}{\gamma} \log(1 + \frac{\gamma}{k(\beta - a_2\alpha)e^{(\beta - a_2\alpha)(x - \frac{y}{a_2})} - \gamma}) + cte; \quad (114)$$

where $k =$ is arbitrary constant.

Case 3.2 : If $a_1 = 0$.

In this situation we takes v as :

$$v = \frac{\partial}{\partial y}, \quad (115)$$

and the invariants are :

$$\chi = x \quad (116)$$

$$\varsigma = u \quad (117)$$

Then $u = \varsigma$, and therefore,

$$u_x = \varsigma_\chi, \quad (118)$$

$$u_y = 0, \quad (119)$$

$$u_{xy} = 0. \quad (120)$$

Substituting these expressions into the Thomas equation, then the solution of the corresponding equation is a constant :

$$\alpha u_x = 0, \implies u = cte. \quad (121)$$

4 Conclusion

In this paper by using the criterion of invariance of the equation under the infinitesimal prolonged infinitesimal generators, we find the most Lie point symmetries group of the Thomas equation. Looking the adjoint representation of the obtained symmetry group on its Lie algebra, we have find the preliminary classification of group-invariants solutions. We have seen that the obtained reduced equation in such case can be transformed on known equation by using an appropriate change of the variables. It is interesting to extend this construction to the supersymmetric Thomas equation, more detail in this question will be given elsewhere[13].

5 Appendix

In this Appendix, we introduce the method for finding the invariants of the group generated by the following infinitesimal generator :

$$v = (a_1 - \gamma x) \frac{\partial}{\partial x} + (a_2 + \gamma y) \frac{\partial}{\partial y} + (\beta x - \alpha y) \frac{\partial}{\partial u}. \quad (122)$$

The invariants are found by integrating the corresponding characteristic system, which is

$$\frac{dx}{a_1 - \gamma x} = \frac{dy}{a_2 + \gamma y} = \frac{du}{\beta x - \alpha y}. \quad (123)$$

The first of these two equations :

$$\frac{dx}{a_1 - \gamma x} = \frac{dy}{a_2 + \gamma y}, \quad (124)$$

is easily solved, the solution are :

$$\frac{1}{\gamma} \log(a_1 - \gamma x) = \frac{1}{\gamma} \log(a_2 + \gamma y) + c. \quad (125)$$

So, one of the invariant is :

$$\chi = (a_1 - \gamma x)(a_2 + \gamma y). \quad (126)$$

Note that χ is a constant for all solution of the characteristic system, so we can replace y by $\frac{\chi}{\gamma(a_1 - \gamma x)} - \frac{a_2}{\gamma}$ before integrating. This leads to the equation :

$$\frac{dx}{a_1 - \gamma x} = \frac{du}{\beta x - \frac{\alpha \chi}{\gamma(a_1 - \gamma x)} + \frac{\alpha a_2}{\gamma}}; \quad (127)$$

therefore :

$$\left\{ \frac{\beta x}{a_1 - \gamma x} - \frac{\alpha \chi}{\gamma(a_1 - \gamma x)^2} + \frac{\alpha a_2}{\gamma(a_1 - \gamma x)} \right\} dx = du. \quad (128)$$

Then the equation has the following solution :

$$\frac{-\beta}{\gamma} x - \frac{(\beta a_1 + \alpha a_2)}{\gamma^2} \log(a_1 - \gamma x) - \frac{\alpha \chi}{\gamma^2(a_1 - \gamma x)} = u + k, \quad (129)$$

for k an arbitrary constant. Finally we have :

$$\varsigma = u + \frac{\beta}{\gamma} x + \frac{(\beta a_1 + \alpha a_2)}{\gamma^2} \log(a_1 - \gamma x) + \frac{\alpha \chi}{\gamma^2(a_1 - \gamma x)}, \quad (130)$$

is the second invariant.

References

- [1] Olver P.J, Application of Lie groups to Differential Equation, Springer, New York 1986.
- [2] Stephani H, Differential equations: Their Solution Using Symmetries, Cambridge University Press, 1989.
- [3] Bluman G.W and Kumei, Symmetries and Differentials Equations, Springer, New York 1989.
- [4] Ibragimov N.H, CRC Handbook of Lie Group Analysis of Differential Equation, CRC Press, Boca Raton, 1986.
- [5] Rodríguez M.A. and Winternitz. P, J.PhysA:Maths.Gen 37(2004) and the references quoted therien.
- [6] Lie S and Engel F, Theories der Tranformation gruppen Vol.39 (Leipzig Teubner) 1890.
- [7] Wong W.T. and Fung P.C.W, Linuo cimento Vol.99B 1987.163.
- [8] Zheng K et al., Physica Scripta Vol.40 1989, 705.
- [9] Sakovich S.Yu, Journal Physiacs A: Maths.and Gen.21 1988,L1123
- [10] Guang-Mei.W et al., Gechoslovok Journal of Physics Vol.52 2002, 749.
- [11] Zhenya Y, Gechoslovok Journal of Physics Vol.53 2003, 298.
- [12] Thomas H.C Journal American Chemistry. Soc.66 1944, 16 64.
- [13] A.Ouhadan and E.H . El Kinani, in preparation.